


## mandarin variety fact sheet

# Daisy



### Origin

California, USA: a hybrid mandarin recently developed by the US Department of Agriculture. It is a cross between Fortune clementine and Fremont mandarin. Imported into Australian quarantine 1991 and first released to selected nurserymen in 1995.


### Type

Early to mid season maturity, seedy.

### Market

Fresh, domestic. Very different in appearance to Imperial mandarin and would enter the market several weeks after the beginning of the Imperial season. May have export potential due to highly coloured skin, good eating qualities and its extended marketing period from on-tree storage. Daisy juice is also highly palatable with a complex flavour.

### Marketing season (estimated)



### Internal quality

Very good flavour with little rag and a sugar:acid ratio which has received favourable comments from preliminary taste testing. Acid levels do not decline significantly as the fruit ages on the tree. This helps to maintain flavour over a long period. The internal characteristics can be explained by the parentage of Daisy. Fremont (clementine x Ponkan mandarin) is an early, sweet mandarin. Fortune (clementine x Dancy mandarin), is a very late, high acid mandarin type and the blending of these two have produced an early/mid season, good quality mandarin. Seed number ranges from 15-20 per fruit. Segments separate easily.

### External quality

Fruit is medium to large (6-8cm in diameter), round with smooth skin. Skin thickness is 4mm and is not difficult to remove. Skin colour is red-orange and almost 'glossy' in appearance. Fruit will hang on the tree for a long period without an obvious deterioration in external appearance. Internal quality begins to deteriorate by mid August in the Sunraysia region.

## Daisy mandarin variety fact sheet

### Postharvest performance

Fruit can be hand 'snapped' from the tree in the early part of its maturity season. The potentially long harvest period and 'robust' nature of the fruit suggests it may have good postharvest handling qualities.

### Field performance

Field grown arboretum trees produced fruit for the first time in 1996. Trees are vigorous, upright, open and thornless. Young trees on trifoliata, citrange, cleopatra and swingle rootstocks are strong and vigorous. The open, upright habit of Daisy would make tree training and shaping much easier than for other mandarin types such as clementines. Grafted Valencia trees on citrange rootstock have shown an alternate bearing habit, whereas trees on trifoliata produced more consistent yields. Crop load manipulation and selective pruning would help to reduce the alternate bearing habit. Fruit was able to tolerate temperatures of 46°C in January 2001 without significant heat damage. Fruit is susceptible to albedo breakdown particularly on Trifoliata rootstock and overwet soil conditions. Soft new flush growth is very sensitive to spray damage and has been 'potch-marked' with zinc, manganese plus oil sprays in spring. Daisy has been included in the South African seed irradiation program to attempt to develop a low seeded Daisy selection. Daisy is also considered self fertile and seed number will not be reduced even if planted in solid, isolated blocks.

### Pest and disease

Susceptible to Emperor Brown Spot (*Alternaria*).

### Extent of plantings

#### Commercial

Currently a very high demand for budwood.

#### Research

Research agency arboreta, small trials and trees planted for budwood supply.

### State of knowledge

very  
limited



very  
high

New variety to Australia that has created commercial interest. It is still being evaluated in South Africa and the USA. It has not been planted extensively in California and budwood was only released to the Florida industry in July 1997.

Growers should ensure that trees are propagated from Approved Budwood obtained from AusCitrus.

**Last Revised:** September 2002

#### Disclaimer:

Information contained in this publication is provided as general advice only. For application to specific circumstances, professional advice should be sought.



NSW Agriculture



Natural Resource  
and Environment

